

On conformally flat initial data for Einstein equations.

Janusz Karkowski and Edward Malec

Institute of Physics, Jagiellonian University, 30-059 Cracow, Reymonta 4, Poland.

We describe conformally flat initial data, with explicitly given analytic extrinsic curvature solving the vacuum momentum constraints. They follow from a solution of Dain and Friedrich discovered in 2001. The cylindrically symmetric subcase of the Bowen-York solution is a subclass of this general configuration.

The process of solving initial constraints of Einstein equations is a difficult task, since they constitute a set of nonlinear partial differential equations. Having said that, it is necessary to point out that these equations are undetermined; thus it comes as an embarrassment that so little is known about their explicit solutions in (at least) some simple cases. The only data that had been available for a long time, up to our knowledge, are those of Bowen-York [1] and (discovered much later) Brandt-Seidel [2], in which vacuum momentum constraints are solved explicitly and it remains to solve only the Lichnerowicz-York equation.

In the first version of this short note we presented another set of explicit solutions of vacuum momentum constraints, obtained by work and guess, that includes a part of the Bowen-York data as a very special subcase. These appear to be included in a solution found by Dain and Friedrich in 2001 [3]. The purpose of this version is to explain the relation between the two solutions.

Let Σ be a Cauchy hypersurface endowed with an internal metric g_{ij} with the scalar curvature R , and the extrinsic curvature K_{ij} . The initial constraint equations in vacuum read [4]

$$\begin{aligned} R &= K_{ij}K^{ij} - (K_i^i)^2 \\ \nabla_i (K_j^i - g_j^i K_l^l) &= 0. \end{aligned} \quad (1)$$

In the case of maximal slicing condition, $K_i^i = 0$, the initial data can be found relatively easily using the conformal method [5]. Let $g_{ij} = \Phi^4 \hat{g}_{ij}$ where $\hat{g}_{ij}$ is the Euclidean metric. The new solution is defined by the following extrinsic curvature, in spherical coordinates,

$$\begin{aligned} K_{23} &= K_{13} = 0 \\ K_1^1 &= \frac{1}{r^3 \Phi^6} \partial_x^2 W \\ K_1^2 &= \frac{1}{\sin \theta r^3 \Phi^6} \partial_r \partial_x W \\ K_2^2 &= \frac{1}{r^2 \sin^2 \theta \Phi^6} \left[\partial_r (r \partial_r W) + \frac{1}{r} (x \partial_x W - W) \right]. \end{aligned} \quad (2)$$

Here $x = \cos \theta$ and W is an arbitrary function of r and θ . It is easy to check that (2) satisfies the momentum constraints, $\nabla_i K_j^i = 0$, where ∇_i is the covariant derivative in the sense of the metric g_{ij} .

This solution generalizes the Bowen-York solution (or, strictly saying, its cylindrically symmetric part with vanishing angular-momentum) of momentum constraints; the latter corresponds to a particular choice of W . For instance, a part of the Bowen-York extrinsic curvature that is displayed below,

$$\begin{aligned} K_{ij} &= \frac{3}{2r^2 \Phi^2} \left(\hat{P}_i \hat{n}_j + \hat{P}_j \hat{n}_i - (\hat{g}_{ij} - \hat{n}_i \hat{n}_j) \hat{P}_l \hat{n}^l \right) + \\ &\quad \frac{3a^2}{2r^4 \Phi^2} \left(\hat{P}_i \hat{n}_j + \hat{P}_j \hat{n}_i + (\hat{g}_{ij} - 5\hat{n}_i \hat{n}_j) \hat{P}_l \hat{n}^l \right), \end{aligned} \quad (3)$$

(with a constant vector $\vec{\hat{P}}$ aligned along the z-axis and $\vec{\hat{n}}$ being a unit normal to centered metric spheres in the $\hat{g}_{ij}$ metric) can be obtained from (2) by substituting

$$W = r \frac{\hat{P}}{2} (x^3 - 3x) \left(1 - \frac{a^2}{r^2} \right). \quad (4)$$

Another special choice of the function W ,

$$W = \frac{\hat{P}}{2} (x^2 + 1), \quad (5)$$

recovers the well known spherically symmetric solution of the momentum constraint

$$\begin{aligned} K_1^1 &= \frac{\hat{P}}{r^3 \Phi^6} = -2K_2^2 = -2K_3^3. \\ K_1^2 &= K_1^3 = K_3^2 = 0. \end{aligned} \quad (6)$$

We would like to point out that (2) is a cylindrically symmetric subset of a result obtained by Dain and Friedrich [3], who found (using the Newman-Penrose edth symbols) the general extrinsic curvature corresponding to the conformally flat initial data. Their results imply ours, if one assumes (in the notation of [3], Sec. 4)

$$W = 2\lambda_2^R(r, \theta) \sin^2 \theta \quad (7)$$

and equates to zero all functions other than λ_2^R in their equations (80 - 82).

There exists yet other set of explicit solutions, in which

$$\begin{aligned} K_{13} &= \frac{1}{r^2} \partial_x Z(r, x) \\ K_{23} &= \frac{1}{\sin \theta} \partial_r Z(r, x) \end{aligned} \quad (8)$$

and the remaining components of K_{ij} do vanish. The function Z does not depend on the ϕ -angle variable, but otherwise is arbitrary. That is the conformally flat version of the Brandt-Seidel solution reported in [2]; its conformally flat generalization can be also found in [3]. A particular form of Z generates the angular-momentum part of the Bowen-York solution. Any linear combination of the two solutions, (2) and (8), obviously solves the momentum constraints.

The remaining hamiltonian constraint reduces to the Lichnerowicz-York equation $\Delta \Phi = -\frac{1}{8} K_{ij} K^{ij} \Phi^5$ (here Δ is the flat laplacian), which determines the conformal factor Φ . One could simulate "multi-black hole" initial data by the appropriate choice of W . The global energy-momentum can be found from standard formulae

$$\begin{aligned} E &= \frac{-1}{2\pi} \int_{S_\infty} d^2 S^i \nabla_i \phi, \\ P_i &= \frac{1}{8\pi} \int_{S_\infty} d^2 S^i K_{ij}. \end{aligned} \quad (9)$$

The asymptotic mass reads $m = \sqrt{E^2 - P_i P^i}$. It is clear that there exists a large set of W 's that generate isolated systems with finite global energy and momentum.

ACKNOWLEDGMENTS

This work was partly supported by the KBN grant 2 PO3B 00623. The authors thank Sergio Dain and Helmut Friedrich for kind information about their results.

-
- [1] J. M. Bowen and J. W. York, *Phys. Rev.* **D21**, 2047(1980).
 - [2] S. R. Brandt and E. Seidel, *Phys. Rev.* **D54**, 1403(1996).
 - [3] S. Dain and Helmut Friedrich, *Commun. Math. Phys.* **222**, 569(2001).
 - [4] C. W. Misner, K. S. Thorne and J. A. Wheeler, *Gravitation*, (H. Freeman and Company, New York, New York, 1973).
 - [5] N. O'Murchadha and J. W. York, *Phys. Rev.* **D10**, 437(1974).